


Drug Receptor Subtypes and Ingestive Behaviour



Neural Mechanisms of ingestive behaviour has long been one of the most intensively investigated areas within neuroscience. Drug studies have been introduced to develop anti-obesity compounds and, more recently, to identify neurotransmitters which might be involved in the control of ingestive behaviour. The current focus within this field is towards neurotransmitter receptors as it has become known that there are multiple receptor subtypes for each identified neurotransmitter. The study of drug receptor subtypes and ingestive behaviour is growing very rapidly, and has become quite complex. This unique volume is designed to guide students and investigators through a number of different neurotransmitter systems and provide them with the latest information on the identities of receptor subtypes most relevant in the study of ingestive behaviour. As such, it will prove enormously useful to all engaged in studies on appetite and obesity treatment. * *

Provides up-to-the-minute reviews of the most exciting new developments in the study of ingestive behavior.* Includes contributions from leading experts in the field.* Tackles the complex issues of relating neurochemical transmission to the neural and behavioral controls of feeding.* The first book of its kind to deal explicitly with drug receptor subtypes in relation to ingestive behavior.

Livros Drug Receptor Subtypes and Ingestive Behaviour - Cooper Serotonin (5-HT) receptor agonists with high affinity for the different subtypes (i.e. and 5-HT₂) receptor families have been shown to affect ingestive behavior. Depressants/pharmacology* Eating/drug effects Feeding Behavior/drug effects **Dopamine Receptor Subtype Agonists and Feeding Behavior** Opioid receptor subtype antagonists and ingestion. In: Cooper SJ, Clifton PG, eds. **Drug Receptor Subtypes and Ingestive Behaviour**. San Diego: Academic **Drug Receptor Subtypes and Ingestive Behaviour (1996-04-17 Drug Receptor Subtypes and Ingestive Behavior (Book) -- Book reviews. Database. Gale Academic OneFile Gale General OneFile Gale Expanded Academic. Effects of selected serotonin 5-HT(1) and 5-HT(2) receptor agonists** Perturbed feeding behaviour and consequential effects on body required for fenfluramine to influence ingestive behaviour (Lucas et al. Another class of drugs that augment serotonin bioavailability are the serotonin

reuptake inhibitors. The established involvement of this particular receptor subtype in **Drug Receptor Subtypes and Ingestive Behaviour**: Neural Mechanisms of ingestive behaviour has long been one of the most intensively investigated areas within neuroscience. Drug studies have been **5-HT(2C) receptor agonists and the control of appetite**. - **NCBI Drug receptor subtypes and ingestive behaviour** / edited by Steven J. Cooper and Peter G. Clifton. Subjects: Receptors, Neurotransmitter. Receptors, Drug. **Drug receptor subtypes and ingestive behaviour** / **University of** Buy Drug Receptor Subtypes and Ingestive Behaviour by Steven J. Cooper, Peter G. Clifton (ISBN: 9780121876203) from Amazons Book Store. Free UK **Neurotransmitters, Drugs and Brain Function - Google Books Result** (1)Kissileff Laboratory for the Study of Human Ingestive Behaviour, School of 5-HT drugs reduce food intake in rodents in a manner consistent with an These effects appear to be mediated, at least in part, by the 5-HT(2C) receptor subtype. **Handbook of Obesity: Etiology and Pathophysiology, Second Edition - Google Books Result** Behavioral Neuroscience, Vol 112(3), Jun 1998, 502-511. .. In S. J. Cooper & P. G. Clifton (Eds.), Drug receptor subtypes and ingestive behaviour (pp. **Drug Receptor Subtypes and Ingestive Behaviour - reviews, first** Integrative Systems and the Development of Anti-Obesity Drugs Tim Kirkham, In Drug Receptor Subtypes and Ingestive Behavior (S. J. Cooper, and P. G. **Serotonin 2C receptor agonists and the behavioural satiety** - **NCBI Neurochemical modulation of ingestive behavior in the ventral pallidum**. A previous report has demonstrated that the blockade of GABAA receptors in the ventral pallidum (VP), a target of Publication Types: D2 Receptor Antagonists Drinking Behavior/drug effects Drug Interactions GABA Agonists/pharmacology **Neural and Metabolic Control of Macronutrient Intake - Google Books Result** opioid peptides and ingestive behavior are reviewed. Role of opiates in feeding behavior: animal studies .. Clifton PG, editors, Drug receptor subtypes. **Naltrexone for the treatment of obesity: review and update - CiteSeerX** To identify the relevant endogenous opioid receptor ligands, mice the tastant, route of drug administration and the specific behavioral test (reviewed in (Bodnar, 1996). receptor subtypes and multiple opioids bind each receptor subtype. To broaden our studies with this aspect of ingestive behavior we **Neurochemical modulation of ingestive behavior in the ventral** Opioid signalling in humans has been implicated in behaviours reinforced by primary .. In: Drug receptor subtypes and ingestive behavior pp. **Opioid Receptor Subtypes and Ingestive Behavior - Contrasting effects of different cannabinoid receptor ligands on mouse ingestive behaviour**. This study characterized the effects of seven diverse cannabinoid receptor agonists (and one antagonist) on ingestive behaviour in Publication Types: Appetite Depressants/pharmacology Appetite Regulation/drug effects* **Book Reviews Drug receptor subtypes and ingestive behaviour** Livros Drug Receptor Subtypes and Ingestive Behaviour - Cooper (0121876209) no Buscape. Compare precos e economize ate 0% comprando agora! : **Drug Receptor Subtypes and Ingestive Behaviour** Neural Mechanisms of ingestive behaviour has long been one of the most intensively investigated areas within neuroscience. Drug studies have been **Drug Receptor Subtypes and Ingestive Behaviour - 1st Edition** Buy Drug Receptor Subtypes and Ingestive Behaviour (1996-04-17) on ? FREE SHIPPING on qualified orders. **DIFFERENTIAL INVOLVEMENT OF ENDOGENOUS OPIOIDS IN** behavior in chicks, and to investigate the interaction with the m1-opioid receptor subtypes in this effect. . In: Drug receptor subtypes and ingestive behavior. **Cannabinoids and the regulation of ingestive behaviour**. - **NCBI Pharmacological differentiation of opioid receptor antagonists by Highly-specific and selective opioid receptor subtype agonists for the mu** (e.g. In: Drug Receptor Subtypes and Ingestive Behavior, (Eds., SJ Cooper and PJ **Drug Receptor Subtypes and Ingestive Behaviour: 9780121876203** Drug receptor subtypes and ingestive behaviour. Edited by SJ Cooper and PG Clifton. Academic Press, New York and London, 1996, 416 pp, ?60.00. ISBN 0 12 **Little and often: Ingestive behavior patterns following hippocampal** Curr Drug Targets. Cannabinoids and the regulation of ingestive behaviour. cannabinoids are mediated through the activation of two receptor subtypes, the Neural Mechanisms of ingestive behaviour has long been one of the most intensively investigated areas within neuroscience. Drug studies have been **Pharmacological differentiation of opioid receptor - Nature** Drug receptor subtypes and ingestive behaviour /. edited by Steven J. Cooper and Peter G. Clifton. imprint. London San Diego : Academic Press, c1996.